

IAS NEWSLETTER



Dr Sachin Tapasvi
President, IAS



Dr SR Sundararajan
General Secretary, IAS
Editor, IAS Newsletter

ARTICLES IN THIS ISSUE

IAS Activities:

IAS CME(Mangalore)(Pg 3-4)

IAS SAARC Sportsmedcon(Kolkata) (Pg 5-6)

IAS Academic Corner:

Expert Talk: Key notes

PCL AVULSION INJURIES: Dr Prateek Kumar Gupta (Pg 7)

Interesting cases:

Interesting Modification to Single Bundle ACL reconstruction to achieve Double Bundle Morphology-Dr Ravinder Kant Manocha (Pg 8-14)

Shoulder instability in elderly: A Different Ball Game-Dr Neha Godghate (Pg 15-18)

IAS office Bearers(2022-2023)



Dr Sachin Tapasvi
President



Dr SR Sundararajan
General Secretary



Dr S Arumugam
Vice President



Dr Sandeep Biraris
Joint Secretary

Governing Council Members



Dr Ashutosh Agrawal



Dr Abhijeet Wahegaonkar



Dr Chirag Thonse



Dr Pradip Nemade



Dr Ramakanth Rajagopalakrishnan



Dr Sagar Kakatkar

Editor- JASSM



Dr Sunit Hazra

Congratulations to the newly elected Office bearers!

Vice President: Dr S Arumugam

EC members: Dr Ramakanth Rajagopalakrishnan


Dr Sagar Kakatkar

Editor: Dr Sunit Hazra

Indian Arthroscopy Society



CME
In Association With



**A.J. Hospital & Research Center,
Mangalore**

As a part of the outreach initiative, IAS conducted the first CME programme on Basics of ACL & Meniscus on 06.03.2022 at A.J Hospital and research centre, Mangalore.






- Dr Sudarshan Bhandary and his team organized an excellent interactive academic programme.
- There were three live surgeries- ACL reconstruction(ACLR) with ITB tenodesis(Dr IPS Oberoi), ACLR with BPTB graft(Dr SR Sundararajan) & ACLR using Quadriceps graft(Dr Nagaraj Shetty) followed by numerous talks by national and local faculty.
- This programme was well received and was attended by more than 50 delegates & was also streamed live on Youtube.

Kudos to team Mangalore for the excellent academic programme!



- The very first SAARC sportsmedcon was held at Kolkata from 25th to 27th March 2022.
- More than 85 international and national faculty participated in this programme to make it a grand success.
- It was a well-attended program with about 150 delegates registered for the conference.
- We had as many as 800 viewers on our YouTube channel as well with viewers from all the SAARC countries.



LIVE SURGERY WORKSHOP
R G Kar Medical College & Hospital
Kolkata

KNEE SURGERIES

- **Medial Meniscus Repair** Supported by Stryker
Dr Anant Joshi
- **ACL** Supported by Conmed
Dr Kanchan Bhattacharyya
- **HTO** Supported by Miraculous
Dr Sachin Tapasvi
- **PCL** Supported by S & N
Dr Roshan Wade, Dr Sandeep Biraris
- **MPFL**
Dr M N Basu Mallick
- **ACL** Supported by MJ Surgicals
Dr Ashish Kumar

SHOULDER SURGERIES

▪ Bankart Supported by Stryker Dr Debasis Chatterjee	▪ Latarjet Supported by Biotek Dr Swarnendu Samanta
▪ Rotator Cuff Supported by Mitek Dr Raju Easwaran	▪ Bankart Supported by Mitek Dr Abheek Kar

- The first day was a day reserved for live surgeries at Rg Kar Hospital. We had the privilege of witnessing as many as nine live surgeries on that single day with Faculty from all over India showcasing their skills for the delegates to grasp the scientific content of the main conference on 26th and 27th March which was also appreciated by delegates.

- Thanks to the IAS team: IPS Oberoi, SR Sundararajan, Sachin Tapasvi, Kolkata Arthroscopy & Sports Surgery Society (KASS) team, Kanchan Bhattacharya, Debasis Chatterjee, Swarnendu Samanta, as well as IAS SAARC Convenor Rajeev Raman for their help & support.
- On behalf of the whole organizing team me Org secretary Dr Sunit Hazra and Scientific Convenor Dr Somudeep Dutta, IAS SAAC Convenor Dr Rajeev Raman would like to thank all delegates and faculty for making this possible on short notice, Long live IAS!!

-By the Organising Team

EXPERT TALK- PCL AVULSION INJURIES




Dr Prateek Gupta
M.S, D.Orth,Mch, FICS(USA), FRCS(UK)
Consultant Orthopedics and Sports surgeon
Sports Medicine Unit, Sir Ganga Ram Hospital
New Delhi

Highlights:

- PCL avulsion injuries are not uncommon injuries
- A high degree of clinical suspicion of instability & stress radiographs are mandatory before proceeding with operative management
- Treatment should be individualized based on size, displacement & comminution of the fragment
- Arthroscopic PCL fixation is a good technique with low morbidity
- Close monitoring & Rehab during the early healing phase is required for achieving optimal outcomes.

INTERESTING MODIFICATION TO SINGLE BUNDLE ACL RECONSTRUCTION TO ACHIEVE DOUBLE BUNDLE MORPHOLOGY



Dr Ravinder Kant Manocha

Northern Railways Central Hospital
Panchkuyian Road, Connaught Place
New Delhi

Dr Maieen-ul Malik

Northern Railways Central Hospital
Panchkuyian Road, Connaught Place
New Delhi

Introduction:

Single Bundle ACL (SBACL) reconstruction has given good results but follow up showed some anterior laxity on Lachman test, pivot glide and reduced rotational stability(1)(2)(3). Most of this deficiency is attributed to failure of SBACL to restore the anatomy and biomechanics of native ACL.

Double Bundle ACL (DBACL) reconstruction, other alternative to better restore anatomy remains controversial. High quality, prospective, randomised studies have found from “no difference”(4) to “better anterior and rotational stability”(5) “better ROM & Objective functional scores”(6) to “fewer revisions”(7). The currently practised technique of Double bundle ACL reconstruction was described in 2004 by Yasuda et al(8).

Much of the controversy regarding four tunnel DBACL advisability as routine alternative to the conventional single-bundle ACL reconstruction(9)(10)(11) is on account of: this being a technically challenging procedure, its pitfalls like longer surgery, abundant fixation, tunnel coalescence with greater complication rate and difficult revision. There are concerns that drilling 4 tunnels for double-bundle ACL reconstructions may lead to problems with bone stock and consequent tunnel enlargement(12)(13). Markolf et al(14) demonstrated that a DBACL reconstruction significantly over-constrained the anterior tibial translation from 30° to 90° of flexion compared with the ACL-intact knee.

A modification that reduces number of tunnels to just Solitary Tibial and Solitary Principal Femoral Socket with a narrow 4mm Supplementary femoral tunnel is suggested to obtain double bundle morphology with essentially a technique similar to Single bundle ACL reconstruction

Review of Literature :

Literature describes reconstruction of Double Bundle Morphology of ACL by number of tunnels varying from two, three and four tunnels. Currently practiced technique, pioneered by Yasuda(9) is a four Tunnel Technique with independent Tunnels in Tibia and Sockets in femur respectively for AM & PL bundles and is named as Anatomic Double bundle reconstruction.

Three tunnel technique for achieving a double bundle morphology was first proposed by Rosengraf(15). These three tunnels can either be two in Femur and one in Tibial or one in femur socket and two in Tibia. Cadaveric studies have been conducted by Petersen et al Mae et al(17), Yagi et al(18). Yagi et al reported that anterior tibial translation was significantly less at full extension and 30° of knee flexion; and under combined rotatory loads of internal tibial torque and valgus torque, the coupled anterior tibial translation was also significantly less at both 15° and 30° of flexion angles.

Two tunnel technique was described by Hemanth et al(19). Two bundle morphology was achieved with solitary tibial and solitary femoral tunnel. Authors demonstrated improved control of knee stability in cadaveric study compared to single tunnel single bundle technique. Authors obtained double bundle morphology by “spreading apart”, two strands of Gracilis+SemiT from remaining two strands of Gracilis+SemiT by a special graft positioning tool and fixed in “spread apart” position by strategically inserting femoral Intrafix® screw between two halves of the graft with diameter of screw determining separation of two bundles.

Surgical technique:

Figure 1A shows the schematic representation of the surgical technique. Steps of Technique are shown in Figure 1B (Graft preparation schematic) & Figure 2 (A to J) and are explained hereinbelow.

1. Graft preparation: 5 Strand graft is prepared with 3 Strands of Semi T and two strands of Gracilis. Gracilis strands are suspended through additional Fibretape loop as shown in Figure 1B.
2. Principal femoral socket is created through AM portal using over the top offset zig by Inside-out Technique to diameter of 5 strand graft. Outlines of Footprint are carefully delineated (Figure 2A).
3. 4mm tunnel is drilled using Outside-in technique. Drill bit is left in situ for parking a shuttling thread (Figure 2B).
4. Tibial Tunnel is drilled by standard outside-in technique (Figure 2C).
5. Two shuttling threads are parked. Shuttling thread of 4mm tunnel is carefully manoeuvred in posterior position (Figure 2D). While pulling out two threads through solitary tibial tunnel, care is taken to prevent entanglement
6. The bone bridge between the principal socket and supplementary tunnel is decorticated to a shallow trough (Figure 2E).
7. All strands of graft are seated into principal tunnel using main shuttling thread, taking care to pull in increments while also simultaneously checking free sliding of second shuttling thread. While the graft is being seated in principal tunnel a Fibre tape loop/No button adjustable loop is also incrementally pulled into narrow tunnel but is kept loose (Figure 2F). Button is flipped as routine.




Figure 1: Schematic representation of the surgical technique. A. Schematic Representation of Technique. Note reconstruction of two bundle morphology. B: Graft is suspended from the Adjustable loop. Fibretape Loop is suspending Gracilis Strands.

8. Fibre tape loop is then pulled tight and secured on lateral cortex of femur with ABS button. Two strands of Gracilis get pulled to the mouth of the narrow tunnel creating two bundle morphology (Figure 2G).

9. Two strands of Gracilis are cycled first and fixed on a post ((Figure 2I = swivel lock anchor) in full extension placing gracilis strands in posterolateral position in tibial tunnel.

10. Three strands of Semi T are cycled and an interference screw is inserted in 30 to 45 Degree flexion between two strands of Gracilis already fixed posterolaterally and Semi T strands held tight antero-medially (Figure 2J). Two parts of the graft thus get separated by the diameter of the interference screw.

11. Immediate post op care remains the standard

12. Rehabilitation protocol is steady and deliberate taking care to avoid any acceleration of rehabilitation

Discussion:

Modification described above is primarily an extrapolation of the idea of Hemanth et al(19) of obtaining double bundle morphology by “spreading apart” strands of graft. As Femoral Intrafix is not available in India; a strategy to achieve “spread apart” was devised, by providing an additional anchorage to Gracilis strands. This strategy is an intermediate between solitary femoral socket DB technique and two femoral sockets technique. Like former, all the stands of the hamstring go into the principal socket and double-bundle morphology is obtained by separating some strands from the remaining strands. Like the latter, there is another tunnel in the femoral condyle albeit a narrow 4mm tunnel. The difference is that no strand actually enters in supplementary tunnel and this tunnel just reroutes two strands of Gracilis from principal tunnel to mouth of narrow tunnel (Figure 2H).

Supplementary tunnel acts merely as additional anchorage point for Gracilis strands. Gracilis courses to tibial tunnel from accessory anchorage point while trebled Semi T courses straight from Principal Socket to tibial tunnel.

Advantages of the modification include achieving two bundle morphology: without making the procedure much longer; making it possible with an intermediate technical expertise with lower complication rate; reduced encroachment of bone stock in short individuals; and easier revision. Second tunnel being narrow and drilled outside-in allows choosing starting point safely away from exit point of principal tunnel of ACL or tunnel of LCL in multi-ligament setting reducing tunnels coalescence risk. Cost of implants is reduced because additional fixation is just a Fibretape loop and suture disc. Post fixation on tibia can easily be 4.5mm cortical screw with washer and can be removed after interference fixation in tibial tunnel is done.

All these features, while achieving double-bundle morphology make this surgery, technically speaking, much closer to a Standard SBACL reconstruction than a Three tunnel or Four tunnel DBACL reconstruction and that is exactly how we view our modification. For this reason, this technique may also reduce the chances of over-constrained anterior translation compared to an ACL intact knee which has been described(14) in Anatomic DBACL technique.

Independent tensioning and fixation of the two bundles at different flexed positions of the knee has either been achieved in four tunnel technique(8) or three tunnel technique with two tibial tunnels(16) and only in cadaveric studies in three tunnel technique with two femoral sockets(17)(18). One of notable merits of our technique is that after fixation of all strands in principal tunnel with suspensory fixation and fixation of gracilis strands to accessory anchorage point, Gracilis strands

are fixed in full extension on a post and then the knee may be moved to a desirable flexed position to fix all the strands in the tunnel. After interference fixation post becomes superfluous as Differential tension and length differential gets built in different strands once interference fixation is done and post can then be safely removed.

“Insertional Site area” concept has helped us understand the place of SB & DB reconstruction in ACL surgeries(20). Siebold published tables detailing the diameters of drill bit and the angle of drilling that should be used for achieving “complete foot print coverage” with SBACL for a given Tibial insertional size. Details of combination of drills sizes, grafts sizes, and angles of drilling were also detailed for achieving complete foot print coverage by DBACL reconstruction assuming a bone bridge of 2mm. He found that DBACL reconstruction for tibial foot print size of less than 14mm grafts would be 5.0 & 5.5mm which are thin and fraught with high risk of graft failure. A minimum acceptable size of 6mm graft diameter for AM bundle was practical with only a Tibial foot print size of 15mm. Considering Femoral footprints tend to be larger by 2mm to tibial foot print, a corresponding minimum acceptable femoral foot print size for DBACL reconstruction would therefore be 17mm. In Indian population, footprints tend to be generally smaller and therefore a small minority of Indian patients may be suitable for Four Tunnel DBACL reconstruction and that is indeed so in our experience. Our modification permits achievement of double bundle morphology of Reconstructed ACL in

much greater proportion of population. There is solitary tibial tunnel, tibial foot print size limiting applicability of our technique due to small tibial foot print becomes non-issue. In effect, short femoral footprint also does not bar any patients from our technique of DBACL because second tunnel doesn't need to be drilled to diameter of PL graft as the graft is not supposed to be insinuated into the supplementary tunnel but held tightly pressed against the mouth of the supplementary tunnel by a fibretape loop. Ordinarily 4mm diameter tunnel is made, but beath pin hole of 2.5mm may be used for supplementary anchorage for two strands of Gracilis. A nitinol wire loop through Meniscus Mender needle can be used for parking second shuttling thread through hole of 2.5mm. Only modification required would be to use Fibrewire instead of fibretape.

A narrow second tunnel doesn't make it mandatory even to leave a minimum of 2mm bony bridge between tunnels. The second tunnel is drilled outside in leaving a safe 4-5mm distance between tunnels at the surface of lateral femoral cortex. On the inside opening the two tunnels may become as much closure as to be right next to each other. Only modification in such situation would be not to create trough in between. Even though the separation between bundles would be small, but as supplementary tunnel is made at the anterior edge of footprint close to cartilage margin, foot print coverage would still be complete. We have achieved two bundle morphology and complete foot print coverage for femoral foot print as small as 9-10mm.

There may be a few pitfalls as well. The obvious would be concern regarding healing of rerouted segment of Gracilis graft on the medial surface of the lateral condyle. Decortication to cancellous surface and gouging out a shallow trough sometimes is an attempt to facilitate such healing. It is our

hypothesis that healing to a flat decorticated cancellous surface may not be a problem as the surface available to Gracilis tendon is increased compared to independent tunnel scenario. It is seated in socket as in any ACL reconstruction, pressed against decorticated cancellous surface (much the same way as rotator cuff tendons pressed against cancellous bone of greater tuberosity in double row RC repair) and finally held to the mouth of 4mm tunnel.

Other pitfalls may be entanglement of Shuttling loops inside tibial Tunnel or fibretape loop getting entwined with graft and getting pulled into principal socket and getting stuck. Loops should be pulled inside tibial tunnel one at a time with suture retriever only, taking care to open prongs of instrument only under vision. The graft should be pulled in small deliberate increments while simultaneously checking to and fro sliding of the fibretape loop, using nylon loop, till the graft is seated and button is flipped. If one still gets stuck, one may just pull out the second shuttling thread or fibretape loop, ignore the second narrow tunnel and complete the procedure as a standard SBACL. However, if technique is followed diligently these complications do not happen. Except in first 2-3 cases when we were still sorting out intricacies of the technique, these complications have not happened in a series of 17 cases.

A more common complication is failure to ensure a strict posterolateral position for Gracilis strands and Anteromedial position for SemiT strands which may happen in about 25% cases.




Figure 2(A-J): Steps of Surgical Technique.

The second shuttle thread is carefully maintained in the posterolateral position and Gracilis strands are fixed first in the posterolateral position to protect against this but accept it if it happens because clinical outcomes of these patients have been as good as in others.

Conclusion:

Described here is a technique which is sort of an intermediate between “Two Tunnel Double bundle” and “Three tunnel Double Bundle technique”. Third Tunnel is a narrow 4mm tunnel which is easily added to a standard SBACL (Two tunnel) technique. Modification can restore ACL morphology much closer to native anatomy than Standard SBACL and has potential to reduce all complications of Conventional Double bundle techniques while restoring double bundle morphology fairly well

References:

1. Tashman S, Collon D, Anderson K, Kolowich P, Anderst W. Abnormal rotational knee motion during running after anterior cruciate ligament reconstruction. *Am J Sports Med.* 2004;32(4):975–983.
2. Yunes M, Richmond JC, Engels EA, Pinczewski LA. Patellar versus hamstring tendons in anterior cruciate ligament reconstruction: a metaanalysis. *Arthroscopy.* 2001;17(3):248–257.
3. Freedman KB, D’Amato MJ, Nedeff DD, Kaz A, Bach BR Jr. Arthroscopic anterior cruciate ligament reconstruction: a metaanalysis comparing patellar tendon and hamstring tendon autografts. *Am J Sports Med.* 2003;
4. Streich NA, Friedrich K, Gotterbarm T, Schmitt H. Reconstruction of the ACL with a semitendinosus tendon graft: a prospective randomized single blinded comparison of double-bundle versus single-bundle technique in male athletes. *Knee Surg Sports Traumatol Arthrosc.* 2008;16:232–238.

5. Muneta T, Koga H, Mochizuki T, Ju YJ, Hara K, Nimura A, Yagishita K, Sekiya I. A prospective randomized study of 4-strand semitendinosus tendon anterior cruciate ligament reconstruction comparing single-bundle and double-bundle techniques. *Arthroscopy* 2007;23:618-628.
6. Marcacci M, Molgora AP, Zaffagnini S, Vascellari A, Iacono F, Presti ML. Anatomic double-bundle anterior cruciate ligament reconstruction with hamstrings. *Arthroscopy* 397 2003;19:540-546.
7. Sally Jarvela, Tommi Kiekara, Piia Suomalainen, Timo Jarvela. Double-Bundle Versus Single-Bundle Anterior Cruciate Ligament Reconstruction: A Prospective Randomized Study With 10-Year Result. *Am J Sports Med.* 2017 Sep;45(11):2578-2585. doi: 10.1177/0363546517712231. Epub 2017 Jun 29.
8. Yasuda K, Kondo E, Ichiyama H, Kitamura N, Tanabe Y, Tohyama H, Minami A. Anatomical reconstruction of the anteromedial and posterolateral bundles of the anterior cruciate ligament using hamstring tendon grafts. *Arthroscopy* 2004;20:1015-1025.
9. Gabriel MT, Wong EK, Woo SL, et al. Distribution of in situ forces in the anterior cruciate ligament in response to rotatory loads. *J Orthop Res* 2004;22:85-9.
10. Cha PS, Brucker PU, West RV, et al. Arthroscopic double-bundle anterior cruciate ligament reconstruction: an anatomic approach. *Arthroscopy* 2005;21: 1275.e1-e8.
11. Vidal AF, Brucker PU, Fu FH. Anatomic double-bundle anterior cruciate ligament reconstruction using tibialis anterior tendon allografts. *Oper Tech Orthop* 2005;15:140-5.
12. Colombet P, Robinson J, Jambou S, Allard M, Bousquet V, de Lavigne C. Two-bundle, four-tunnel anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2006;14(7): 629-636.
13. Siebold R. Observations on bone tunnel enlargement after double bundle anterior cruciate ligament reconstruction. *Arthroscopy.* 2007;23(3):291-298.
14. Markolf KL, Park S, Jackson SR, McAllister DR. Anterior-posterior and rotatory stability of single and double-bundle anterior cruciate ligament reconstructions. *J Bone Joint Surg Am.* 2009; 91(1):107-118. [PubMed: 19122085]
15. Rosenberg TD, Graf B. Techniques for ACL reconstruction with Multi-Trac drill guide. Mansfield, MA: Acufex Microsurgical Inc, 1994
16. Petersen W, Tretow H, Weimann A, Herbort M, Fu FH, Raschke M, Zantop T. Biomechanical evaluation of two techniques for double-bundle anterior cruciate ligament reconstruction: One tibial tunnel versus two tibial tunnels. *Am J Sports Med* 2007;35:228-234.
17. Mae T, Shino K, Miyama T, Shinjo H, Ochi T, Yoshikawa H, Fujie H. Single- versus two-femoral socket anterior cruciate ligament reconstruction technique: Biomechanical analysis using a robotic simulator. *Arthroscopy* 2001;17:708-716
18. Yagi M, Wong EK, Kanamori A, Debski RE, Fu FH, Woo SL. Biomechanical analysis of an anatomic anterior cruciate ligament reconstruction. *Am J Sports Med* 2002;30:660-666.
19. Hemanth R, Gadikota MS, Jia-Lin Wu. Single-tunnel double-bundle anterior cruciate ligament reconstruction with anatomical placement of hamstring tendon graft: can it restore normal knee joint kinematics? *Am J Sports Med* 2010;38:713-2012.
20. Siebold R The concept of complete footprint restoration with guidelines for single- and double-bundle ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* (2011) 19:699-706 DOI 10.1007/s00167-010-1376-x

SHOULDER INSTABILITY IN ELDERLY: A DIFFERENT BALL GAME



Dr. Neha Godghate
MS(Ortho), MCh(UK), DNB
Consultant, Grace Orthocare
Hospital, Nagpur, Maharashtra

Dr. Ninad Godghate
MS(Ortho), MCh(UK).
Grace Orthocare Hospital,
Nagpur, Maharashtra

Dr. Krishnamohan Saindane
MS(Ortho), FICS.
Grace Orthocare Hospital,
Nagpur, Maharashtra

Abstract:

Shoulder instability in elderly is relatively uncommon and less studied condition. We report a case of shoulder instability in a 75 years old active elderly independent lady.

Even after conservative management her symptoms persisted. On investigations, she was diagnosed to have full thickness Supraspinatus tear with Bankart's lesion. The repair of both Supraspinatus and Bankart's lesion was done arthroscopically. At the end of 5 months, she was able to do all her pre-fall activities without any apprehension.

Introduction:

Though shoulder instability is commonly witnessed in younger individuals, the incidence of instability is increasing in middle aged and elderly population as well(1). The reason being increasing life expectancy and increasing participation in leisure sports in elderly. In elderly, anterior dislocation of shoulder is the commonest like young dislocators. However, the pathophysiology differs(2). In contrast to young individuals, a minor force is required to dislocate the shoulder in elderly. Age-dependent pathophysiology of anterior shoulder dislocation is attributable to changes in tissue elasticity with age.(3). As a result of various biological changes, we get to see capsular lesions commonly than labral tears.(4). Neurovascular injuries are more common, especially of axillary nerve. The incidence of nerve injury is upto 9.3%,(5) and deLaat et al.(6) calculated the prevalence of nerve injury as follows, axillary nerve (37%), supra scapular nerve (29%), radial nerve (22%). However, axillary nerve injuries recover between 6 weeks and a year after dislocation (6,7). Few studies have demonstrated the incidence of

dislocation in old population (above 60 years) is upto 20%-44%(8,9,5,10) and recurrence rate ranges from 10% to 31%. (5,8,11,12,13,14)

Case Report:

A 75 years old right hand dominant lady presented to OPD with history of fall at home a day prior, when she felt her right shoulder suddenly popping out and going in. She presented with main complaints of feeling of instability and pain. Patient had no previous history of shoulder pain. On examination, Supraspinatus strength was grade 4, Full-can test was positive. Apprehension test was positive. Neurological examination was within normal limits. On day 1, there was no evidence of any bony injuries or arthritis on X-Ray. She was conservatively managed for 3 weeks with medications and sling support. At the end of 3 weeks her symptoms and clinical findings persisted. MRI of right shoulder was done which showed typical Bankart's lesion with a full thickness rotator cuff tear(Fig 1). It was decided to repair both the lesions arthroscopically.

Technique:

In lateral decubitus position, a diagnostic round of glenohumeral joint showed fraying of biceps. Hence, biceps tenotomy was performed. Glenoid labrum was liberated and shaving of the frayed edges was done. Rasp was used to abrade the medial edge of glenoid for better healing. Two 2.5 mm anchors were used for Bankart's repair(Fig 2A& 2B). After achieving robust soft tissue bumper, arthroscope was shifted to subacromial space. After a thorough subacromial decompression, one 5.5 mm double loaded suture anchor was inserted. With the help of ante-grade suture passer, sutures were passed through the cuff tissue. Sutures were then tied with sliding knots. A watertight repair achieved(Fig 2C & 2D). Post-operatively, the patient was immobilised in an immobiliser. Pendulum exercises were started from day 2. Gradual range of motion exercises were started from 2nd week. Strengthening exercises were started from 6th week.




Fig 1: Pre-operative MR imaging. A: Axial cuts showing labral tear, B: Coronal cuts showing a rotator cuff tear




Fig 2: Arthroscopic images of labral repair & cuff repair. A&B: Labral repair. C&D: Rotator cuff repair

Results:

At the end of 2.5 months Range of motion improved gradually. At the end 5.5 months, VAS score was significantly reduced from 9 (pre-operative) to 2. Constant score (15) improved to 81 at the end of 5.5 months (Fig 3).

Discussion:

As far as shoulder instability is concerned, rotator cuff and ligamentous structures act in synergism. Rotator cuff provides significant dynamic stability to shoulder via “concavity compression mechanism”. Hence, loss of cuff integrity jeopardises stability of glenohumeral joint. The injury to anterior support which was traditionally thought to be insignificant in elderly has now shown to be statistically significant factor in maintaining stability.(14) Anterior labral injuries must not be overlooked in elderly patients with anterior shoulder dislocations. In one study every patient with anterior dislocation were associated with an anterior labral injury (5), and in another a concomitant large

or massive tear of the rotator cuff, a concomitant Bankart lesion, or an anterior glenoid rim fracture were shown to be causative factors for redislocation in patients over the age of 55 (13) .

Recurrent dislocations in the elderly is not as rare as once thought. Various studies have reported the recurrence rate ranging from 10%- to 31%.(5,8,11,12,13,14). Despite the extensive literature on surgical management of shoulder instability in young individuals, there is a paucity of literature regarding outcomes in older. Controversy existed in management. Shin et al. has given an algorithm to address this dilemma(1). Levy et al. have suggested that multiple recurrences are due to disruption of both anterior and posterior stability mechanisms(13). The studies by Voos et al. and Shield et al. showed that after repair of both the structures, functional and clinical outcomes of affected and unaffected extremity were similar at the end of 2.7 years and 3 years respectively(16,17). The largest cohort study was done in 2020 by Chan et al. in which 103 middle-aged and elderly patients with



Fig 3: Clinical follow up at 5.5 months. A: Full Range of movement B: No apprehension

shoulder instability were treated surgically. They concluded that none of the patients had any instability event at 2 years follow up(18)

Conclusion:

As elderly patients remain active longer into life, orthopaedic surgeons specializing in sports medicine may see more patients with combined rotator cuff and labral injuries after shoulder dislocations. In case of shoulder instability in the elderly, we need to address both anterior as well as posterior stabilising mechanisms for better functional outcomes. Further studies are needed to facilitate more accurate patient counselling with regards to outcome expectations after treatment.

References:

1. Shin SJ, Young Won-ko, Jueob Lee. Traumatic anterior shoulder dislocation in elderly patients. *Arthrosc Orthop Sports Med* 2016;3(1):24-31
2. De Beer J, Bhatia D. Shoulder instability in middle aged and elderly patients: Pathology and surgical implications 87 *International Journal of Shoulder Surgery* - Oct-Dec 2010 / Vol 4 / Issue 4
3. Maier M, Geiger EV, Ilius C, Frank J, Marzi I. Midterm results after operatively stabilised shoulder dislocations in elderly patients. *Int Orthop* 2009;33:719-23.
4. Mizuno N, Yoneda, Nakagawa, Hayashida. The pathology of anterior capsule in patients over 40 years with recurrent shoulder dislocation *Int. Orthop. (SICOT)*
5. Gumina S, Postacchini F. Anterior dislocation of the shoulder in elderly patients. *J Bone Joint Surg Br* 1997;79:540-3.
6. de Laat EA, Visser CP, Coene LN, Pahlplatz PV, Tavy DL. Nerve lesions in primary shoulder dislocations and humeral neck fractures. A prospective clinical and EMG study. *J Bone Joint Surg Br* 1994;76:381-3.
7. Hawkins RJ, Bell RH, Hawkins RH, Koppert GJ. Anterior dislocation of the shoulder in the older patient. *Clin Orthop Relat Res* 1986;(206):192-5.
8. Rowe CR. Prognosis in dislocations of the shoulder. *J Bone Joint Surg Am* 1956;38:957-77.
9. Kazár B, Relovszky E. Prognosis of primary dislocation of the shoulder. *Acta Orthop Scand* 1969;40:216-24.
10. Murthi et al. Shoulder dislocation in older patient. *JAAOS* 2012
11. Stayner et al. Shoulder dislocation in patients older than 40 years of age *OCNA* 2000
12. Neviasser RJ, Neviasser TJ, Neviasser JS. Concurrent rupture of the rotator cuff and anterior dislocation of the shoulder in the older patient. *J Bone Joint Surg Am* 1988;70:1308-11.
13. Levy O, Pritsch M, Rath E. An operative technique for recurrent shoulder dislocations in older patients. *J Shoulder Elbow Surg* 1999;8:452-7.
14. Rapariz JM, Martin-Martin S, Pareja-Bezares A, Ortega-Klein J. Shoulder dislocation in patients older than 60 years of age. *Int J Shoulder Surg* 2010;4:88-92.
15. Constant CR, Murley AH. A clinical method of functional assessment of the shoulder. *Clin Orthop Relat Res.* 1987 Jan; (214):160-4. PMID: 3791738.
16. Voos, James E., et al. "Outcomes of combined arthroscopic rotator cuff and labral repair." *The American Journal of Sports Medicine* 35.7 (2007): 1174-1179.
17. Shields, Edward, et al. "Functional and imaging outcomes of arthroscopic simultaneous rotator cuff repair and Bankart repair after shoulder dislocations." *The American Journal of Sports Medicine* 42.11 (2014): 2614-2620.
18. Chan Wayne W. Chan, Tyler J. Brolin, Ocean Thakar, Manan S. Patel, Daniel S. Sholder, Joseph A. Abboud, Charles L. Getz. Concomitant rotator cuff repair and instability surgery provide good patient-reported functional outcomes in patients aged 40 years or older with shoulder dislocation. *JSES International*, Volume 4, Issue 4, 2020, 792-796